

Nonequilibrium charge density wave ordering from anomalous velocity in itinerant helical magnets

Cenke Xu¹ and J. E. Moore^{1,2}

¹*Department of Physics, University of California, Berkeley, CA 94720*

²*Materials Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720*

(Dated: February 2, 2008)

The Karplus-Luttinger anomalous velocity is shown to lead to electric-field-induced charge accumulation in nearly ferromagnetic noncollinear magnets with itinerant electrons, like *MnSi*. For helical magnetic ordering, the balance between this accumulation and the Coulomb interaction leads to a nonequilibrium charge density wave state with the period of the helix, even when such accumulation is forbidden by an approximate gauge-like symmetry in the absence of electric field. We compute the strength of such charge accumulation as an example of how unexpected many-electron physics is induced by the inclusion of the one-electron Karplus-Luttinger term whenever the local exchange field felt by conduction electrons does not satisfy the current-free Maxwell equations.

PACS numbers: 72.15.Gd 71.45.Lr

Recently the Karplus-Luttinger “anomalous velocity” in crystal structures without inversion symmetry has been the subject of renewed interest. It has been reinterpreted as an additional term in the semiclassical equations of motion resulting from a Berry phase in momentum space¹, and shown to be determined solely by electronic properties at the Fermi surface², consistent with the spirit of Landau’s Fermi liquid theory. A natural question is whether the existence of a new term in the single-electron equation of motion modifies standard interacting electron physics. We answer this question in the affirmative for long-period itinerant helimagnets like *MnSi* and related systems: the anomalous velocity leads to a nonequilibrium steady-state charge density wave (CDW) transverse to an applied electrical field.

Such ordering is forbidden by symmetry in the absence of an electrical field, but the combination of an applied spatially constant electrical field with the spatially variable effective “Hund’s rule” magnetic field yields a charge density wave whose amplitude is balanced by the Coulomb interaction. Similar noncollinear orderings may exist in other systems like dilute magnetic semiconductors, where simple estimates predict a much stronger version of the effect we find. We show that the same type of nonequilibrium charge ordering is generic to noncollinear magnets that support effective magnetic fields (from exchange interactions) that are inconsistent with the current-free Maxwell equations.

As in the two-dimensional quantized Hall effect, there are finite transverse conductances in three-dimensional transport that do not depend on relaxation and hence are “intrinsic.” Here we focus on the Karplus-Luttinger anomalous velocity in magnetic materials with broken time-reversal symmetry (“*T*-breaking materials”), which has been argued to explain the anomalous Hall effect in ferromagnetic semiconductors¹. A similar intrinsic transverse conductance underlies the “spin Hall effect” predicted for some doped semiconductors without broken time-reversal symmetry^{3,4,5}. Current in all these situations flows perpendicular to electric field and another

vector: in the ordinary Hall effect, the other vector is magnetic field; in the anomalous Hall effect, it is the local magnetic moment; and in the proposed intrinsic spin Hall effect, it is the spin polarization that determines the flow of the current. Because the direction of the current is determined by this third-party vector, it is interesting to ask what will happen if this vector is not homogenous.

The standard derivation of the anomalous Hall effect contribution from the Karplus-Luttinger term depends on nearly constant local moments, so the nearly ferromagnetic noncollinear magnets like *MnSi*^{6,7} are natural candidates. An intuitive understanding of the induced CDW we find in such materials can be gained by considering a stacking of several blocks of an itinerant ferromagnet (e.g., iron), in such a way that the local ferromagnetic direction rotates in a helix (Fig. 1). Then under an applied electric field perpendicular to the helix direction, the different Hall currents in each block will lead to a buildup of charge at the junctions between different blocks. The result we find for itinerant helimagnets can be viewed as a continuum version of the above physics, in which the transverse current flow and charge buildup is dominated by the anomalous Hall effect in materials with broken time-reversal and inversion symmetries. This charge buildup will be largely compensated by screening by the conduction electrons, but there is a residual lattice distortion and charge oscillation in the helix direction that should be experimentally detectable.

Long-period modulations (periodicity over several hundred angstroms) are also found in *SrFeO₃* and *CaFeO₃*⁸. We find that in these helimagnets and some other configurations, an external electric field in certain directions will induce a charge density wave, and that this nonequilibrium charge ordering is directly related to the symmetry of the magnetization. The estimates we give of the strength of this charge density wave are easily carried over to other materials; as the anomalous velocity only depends on the electrons or holes close to the Fermi surface², the detailed band structure is irrelevant.

Since we will consider materials with no symmetry under spatial inversion and with nonzero spin-orbit coupling, the leading spin-orbit coupling should be linear in electron momentum. The free-electron Hamiltonian is

$$H = \sum_{\mathbf{k}, \sigma} c_{\mathbf{k}\sigma}^\dagger (\epsilon_{\mathbf{k}} + \gamma \mathbf{k} \cdot \boldsymbol{\sigma}) c_{\mathbf{k}\sigma}. \quad (1)$$

For simplicity, we will later approximate the dispersion relation by $\epsilon_{\mathbf{k}} = \hbar^2/(2m_e) \cdot k^2$. The detailed structure of the spin-orbit coupling term is not too important for the following calculation, and in fact the same effect appears for the Luttinger Hamiltonian for a spin-3/2 band with only $(\mathbf{k} \cdot \mathbf{s})^2$ terms. The modified semiclassical equation of motion is

$$\begin{aligned} \hbar \dot{\mathbf{k}} &= -eE - e\dot{\mathbf{x}} \times \mathbf{B} \\ \dot{\mathbf{x}} &= \frac{\partial \epsilon_{\mathbf{k}}}{\partial \mathbf{k}} - \dot{\mathbf{k}} \times \boldsymbol{\Omega}. \end{aligned} \quad (2)$$

The Berry curvature $\boldsymbol{\Omega}$ can be regarded as a magnetic field in momentum space: in terms of the periodic part u_n of the wavefunction for Bloch band n , its form is

$$\Omega_i(n, \mathbf{k}) = \epsilon_{ijk} \text{Im} \left\langle \frac{\partial u_n}{\partial k_j} \left| \frac{\partial u_n}{\partial k_x} \right. \right\rangle. \quad (3)$$

We can easily see in (1) that the helicity is a conserved quantity, and for the same momentum, electrons with different helicities have different energy, so actually here we have two bands, with two different helicities. These two bands have a degenerate point at $\mathbf{k} = 0$, close to this point, using (3), we will see the Berry phase looks like a monopole in momentum space,

$$\Omega_i(\lambda, k) = \lambda \frac{k_i}{k^3}. \quad (4)$$

Here λ is the helicity $\hat{\mathbf{k}} \cdot \boldsymbol{\sigma}$. Although we use this form for the Berry phase in our calculations, the specific form of the Berry phase is not important: the anomalous Hall effect is basically a Fermi surface effect. If there is a gap between two bands, an infrared cut-off at $k = 0$ is necessary, but this will not contribute to the final result.

Under an electric field $\mathbf{E}$, the solution of (2) has both normal acceleration along electric field and an anomalous velocity in the plane normal to $\mathbf{E}$, leading to an anomalous Hall current¹

$$\mathbf{J} = \sum_{\lambda} \frac{e^2}{\hbar} \int d^3k (\boldsymbol{\Omega}(\lambda) \times \mathbf{E}) f(\varepsilon). \quad (5)$$

Here $f(\varepsilon)$ is the distribution of particles. If $f(\varepsilon)$ is just the equilibrium distribution, the anomalous current is zero because the equilibrium distribution is even under $\mathbf{k} \rightarrow -\mathbf{k}$, and the Berry phase is odd. More generally, the anomalous Hall effect is forbidden by time-reversal symmetry: electric field is even under time reversal, but current is odd, and because the current is perpendicular to electric field, this anomalous Hall current has no dissipation, so without another vector which is odd under

time-reversal there can be no Hall current. In magnetic materials the ordered moments provide this additional vector, analogous to an external applied field.

The ordered moments produce the Hall current via the “local Hund’s rule” coupling to spin, whose origin is in exchange interactions rather than magnetic dipolar interactions. This interaction between itinerant electrons or holes and the local magnetic order can be modelled in a simple Hamiltonian

$$H' = h \mathbf{m} \cdot \mathbf{s}. \quad (6)$$

Here $\mathbf{m}$ is the local magnetic moment. It is assumed that the background moments are strongly ordered so that it is appropriate to use the expectation value of $\mathbf{m}$ as the Hund’s rule field and ignore fluctuations. We will assume for now that the magnetic configuration has constant magnitude and normalize $|\mathbf{m}|^2 = 1$.

The conduction electron spin density $\mathbf{s}$ includes spin-1/2 matrices in MnSi and spin-3/2 for the Luttinger model of GaAs (here we define $\mathbf{s}$ to be dimensionless so that h has units of energy). The exchange field h is proportional to the magnitude of the ordered local moment. If Mn spins are fully polarized, $h = N_{Mn} S J^1$, where N_{Mn} is the density of Mn ions, S is the spin of Mn ion, which is 5/2, and J is the exchange energy between local moment and itinerant electrons, which we take to be 50 meV nm³⁹. In MnSi, the Mn ions are not fully polarized and the local moment is about $0.4\mu_B$ per ion¹⁰. So in this material h is suppressed to 8% of its maximum value. In MnSi, the lattice constant is 4.6 angstroms and there are 4 Mn ions per unit cell, so we obtain $N_{Mn} \simeq 30/\text{nm}^3$. Collecting all the results above, we finally estimate $h = 300$ meV in MnSi.

On symmetry grounds, if both time-reversal and inversion symmetries are broken, one expects a Hall current from the anomalous velocity term:

$$\mathbf{J}_{AH} = \sigma_{AH} \mathbf{E} \times \mathbf{m}. \quad (7)$$

Both sides of this equation are odd under time reversal, so a nonzero anomalous Hall effect is consistent with broken time-reversal and inversion symmetries.

In this problem we have three different energy scales: the Fermi energy, spin-orbit coupling energy, and the Hund’s rule coupling between itinerant spin and local moment. For convenience we assume $E_f \gg \lambda \gamma k_f \gg \lambda h$, although our main result does not depend on this assumption. In this case the basic band structure of this material is still applicable and we can regard the effect of local moment as a perturbation. Suppose that $\mathbf{m}$ is along the $\hat{z}$ direction. Then the Fermi sea with positive helicity will be pushed down along $-\hat{z}$, because if an electron with momentum along negative $\hat{z}$ has positive helicity, it will benefit from the interaction with local moment. For the same reason, the Fermi sea with negative helicity will be pushed up along $\hat{z}$. Thus the Fermi surface has an anisotropic Fermi wavenumber k_F .

Because the energy at the Fermi surface must be constant, we have

$$\frac{\hbar^2}{2m_e}k_f^2 + \lambda\frac{\gamma}{2}k_f = \frac{\hbar^2}{2m_e}k_\theta^2 + \lambda\frac{\gamma}{2}k_\theta + \lambda\frac{h}{2}\cos\theta. \quad (8)$$

Solving these equations, we obtain

$$k_\theta = k_f - \frac{hm_e\lambda\cos\theta}{\hbar^2k_f}. \quad (9)$$

We have used the assumption that the Fermi energy is much greater than the spin-orbit coupling to ignore the γ in the denominator. So the unknown value of γ is irrelevant as long as it satisfies the assumption above. This is analogous to the situation in the Rashba model, which in the absence of interactions and disorder has a universal spin-Hall conductivity independent of spin-orbit coupling⁴. Here k_f is the Fermi vector at $\theta = \pi/2$, m_e is the electron effective mass, θ is the angle between $\mathbf{k}$ and $\mathbf{m}$, and λ is the helicity. Now because of this anisotropy, integrating over the Fermi sea as in (5) and summing over the two helicities finally obtains a nonzero Hall current: the transverse conductivity is

$$\sigma_{AH} \simeq \frac{e^2\hbar m_e}{\hbar^3} \frac{1}{k_f}. \quad (10)$$

For a numerical estimate, take $1/k_f \approx 1 \text{ \AA}$ and replace the effective mass by the bare mass: then the Hall conductivity $\sigma_{AH} \simeq 2 \times 10^3 \Omega^{-1} \text{cm}^{-1}$. In very clean samples of MnSi, the diagonal conductivity σ is about $10 \mu\Omega^{-1} \text{cm}^{-1}$ below 20 K, the temperature below which the local moments exist⁶, so $\sigma_{AH}/\sigma \approx 10^{-4}$. As discussed below, this ratio will increase in slightly disordered samples.

This $1/k_f$ behavior appears strange because it predicts that as the particle density vanishes, the Hall effect will be infinitely large. However, the above calculation is based on the assumption that the coupling with local moment is much weaker than the Fermi energy. If k_f is very small, the Hund's rule coupling dominates, almost all the spins are aligned along the direction of local moment, and it is the spin-orbit coupling that is a perturbation.

All the calculations above are based on the nonvanishing Berry phase, which depends on adiabaticity of electron motion. Heuristically, because $\Delta t \cdot \Delta E \geq \hbar/2$, the relaxation time Δt should be large in order to make sure ΔE is very small compared with the band gap at Fermi surface. Although the precise strength of spin-orbit coupling in the materials we study is unknown, it has to be much bigger than $\hbar$, the coupling to local moment, for the perturbative calculation above to be valid. So if $\tau \gg \hbar/h \approx 10^{-14} \text{s}$, the calculation should be valid: this is a typical relaxation time in a metal in room temperature, so at low temperature our assumptions are satisfied.

Now consider what will happen if the local moment is not perfectly collinear. First we discuss the helical case which is relevant to MnSi and other materials discussed in the introduction⁶⁸. For simplicity, we take local moments

$$\mathbf{m}(x) = \cos(qx)\hat{z} + \sin(qx)\hat{y}. \quad (11)$$




FIG. 1: Geometry of charge accumulation in a helical magnetic background. In an electric field $\mathbf{E}$ perpendicular to the helix direction, a transverse CDW forms with the same modulation vector $\mathbf{q}$ as the helix.

The modulation period $2\pi/q$ is typically at least $100 \text{ \AA}$, which means the local moment is rotating very slowly when we move along $\hat{x}$. Because it is rotating very slowly, particles in different regions of the material only see a homogenous local moment, and locally have the same momentum distribution as previous several paragraphs. First keep the electric field along $\hat{y}$: then based on (7) and (10), the local Hall current density is

$$J_{AH}(x) = \cos(qx)\sigma_{AH}E_y. \quad (12)$$

However, electric charge is always a conserved quantity, and obviously, this anomalous charge current does not satisfy the static continuity condition $\nabla \cdot \mathbf{J}_{AH} = 0$, so charge will accumulate, inducing an electric field. Finally, given some dissipation mechanism, the system will reach a steady state in which the diagonal current from the induced electric field cancels the original Hall current:

$$\sigma E' = -J_{AH}. \quad (13)$$

Here σ is the normal diagonal conductivity of the material. From the Maxwell equation $\nabla \cdot \mathbf{D} = 4\pi\rho$, we obtain the relation between accumulation and electric field

$$\rho(x) = q \sin(qx) \frac{4\pi\epsilon(q)\sigma_{AH}}{\sigma} E_y. \quad (14)$$

$\epsilon(q)$ is the static dielectric function at wave vector q . This $\rho(x)$ describes the **bare** or external electric charge density, which will be strongly screened if $\epsilon(q) \gg 1$.

Let us estimate the magnitude of this charge density wave effect. First, in metal $\epsilon(q)$ is given by $\epsilon(q) = 1 + q_{TF}^2/q^2$, if $q^{-1} = 10 \text{ nm}$, we get $\epsilon(q) \simeq 2 \times 10^4$. Also we have $\sigma_{AH}/\sigma \simeq 10^{-4}$. If we take electric field to be 100 V cm^{-1} , the density modulation is about 10^{14} cm^{-3} , which because of its periodicity should be measureable in a diffraction experiment. Furthermore, from (14) the density modulation is proportional to the ratio between anomalous Hall conductivity and diagonal conductivity. Reducing the diagonal conductivity by adding impurities will lead to increased charge modulation until the relaxation time becomes so short that the Berry's phase assumption discussed above is violated.

This charge density wave exists for helical direction $\hat{x}$ when the electric field has in any component in the yz plane. However, if the field is along $\hat{x}$, the Hall current

will be in yz plane, and it is homogenous in this plane, so $\nabla \cdot \mathbf{J} = 0$ and there is no accumulation. Actually, the formation of charge density wave is because of the breaking of an approximate spiral symmetry possessed by the long-period helical spin state. A homogeneous background has 3 translational symmetries, T_x , T_y and T_z . When the helical magnetic moment is added, it seems that T_x has been broken. However, the system still has a special helical or “screw” symmetry: combine the translation along $\hat{x}$ with a rotation, by defining

$$T'_x = T_x(a)R_x(aq), \quad (15)$$

where a is the lattice constant and $R_x(\theta)$ is the rotation transformation around axis $\hat{x}$. We can see the system is invariant under T'_x , T_y and T_z , and these three operators all commute. When we turn on the electric field in yz plane, this spiral translation symmetry is broken by the external field, and we have a charge density modulation which is forbidden if the helical symmetry is unbroken., and the period is the same with the spiral moment configuration. However, if the electric field is along $\hat{x}$, this spiral symmetry is not broken as long as the external field is homogenous. We can conclude that, although our calculation above was based on an assumption about the three energy scales in the problem, this CDW effect should be relatively independent of these assumptions, because it is controlled by symmetry breaking.

This absence of accumulation in the absence of an applied electric field is strictly correct in the continuum limit where the helical transformations are exact symmetries: lattice effects can lead to a very small accumulation if the helix vector is commensurate with the periodicity of the lattice. In the continuum, charge accumulation is always weak as long as the magnitude of the magnetic ordering is constant, because of a local symmetry.

This absence of accumulation in the absence of an applied electric field is strictly correct in the continuum limit where the helical transformations are exact symmetries: lattice effects can lead to a very small accumulation if the helix vector is commensurate with the periodicity of the lattice. In the continuum, charge accumulation is always weak as long as the magnitude of the magnetic ordering is constant, because of a local symmetry. First write (1) and (6) in operator form:

$$H = \psi^\dagger_\alpha \left(-\frac{\hbar^2}{2m_e} \nabla^2 \delta_{\alpha\beta} - \gamma(i\nabla \cdot \mathbf{s})_{\alpha\beta} \right) \psi_\beta + \hbar \mathbf{m}(\vec{r}) \cdot (\psi^\dagger_\alpha \mathbf{s}_{\alpha\beta} \psi_\beta). \quad (16)$$

Now in this Hamiltonian let the local magnetization $\mathbf{m}$ vary slowly from point to point. The direction of local moments in the whole material can be made uniform by performing a position-dependent rotation, written as

$$R(\vec{r})_{ij} m(\vec{r})_j = m(0)_i. \quad (17)$$

We can keep the interaction H' invariant if we perform another position-dependent $SU(2)$ transformation $U(\vec{r})$

on the spinor ψ_α , as long as

$$U(\vec{r})^\dagger S_i U(\vec{r}) = R(\vec{r})_{ij} S_j. \quad (18)$$

Normally, this kind of symmetry is broken by the kinetic term because the derivative will be modified to $\partial_i \rightarrow \partial_i - U^\dagger \partial_i U$ after a position dependent transformation.

However, because we assumed the variation of local direction is slow, both R_{ij} and U are very slowly varying functions, so we can ignore $U^\dagger \partial_i U$ in the kinetic term. This is equivalent to ignoring derivatives of the $SU(2)$ transformation. Because the AHE only involves particles close to the Fermi surface, as long as the characteristic length scale of variation of local moment is much larger than $1/k_f$, the derivatives $U^\dagger \partial_i U$ do not modify the spectrum close to the Fermi surface and can be neglected.

Because of this local $SU(2)$ symmetry, the slowly varying system is equivalent to the case with homogenous magnetization, and without electric field the charges are homogeneously distributed. However, when the magnitude of local moment varies, the $SU(2)$ gauge symmetry is broken as we can never perform any position dependent rotation to transform this system to the homogenous magnetization case. When we add certain external fields to the system with constant $|\mathbf{m}|$, the gauge symmetry is also broken: local rotation transforms the external field, and without protection by symmetry, there will generically be an inhomogenous charge distribution.

In the helical case, an electric field along the helix direction preserves the symmetry and the charge density remains homogenous. It is interesting to look for magnetic moment configurations in which the charges remain homogenous for any applied electric field. The constant magnetic moment is certainly an example, but there are others: to prevent charge accumulation, the Hall current must satisfy $\nabla \cdot \mathbf{J}_{AH} = 0$. This gives a constraint on the moment configuration:

$$\begin{aligned} \nabla \cdot \mathbf{J}_{AH} &= \sigma_{AH} \nabla \cdot (\mathbf{m} \times \mathbf{E}) \\ &= \sigma_{AH} (\mathbf{E} \cdot \nabla \times \mathbf{m} - \mathbf{m} \cdot \nabla \times \mathbf{E}). \end{aligned} \quad (19)$$

The second term in the bracket of the second line is zero because curl of electric field is always zero. Hence there is no charge accumulation in any electric field if $\mathbf{m}$ has zero curl. This requirement is satisfied if $\mathbf{m} = \nabla \phi$, so the moment configurations with no accumulation are equivalent to source-free *electric* field configurations of constant magnitude. One example is the monopole configuration, in which $\mathbf{m}(\mathbf{r}) = \hat{r}$ with the potential $\phi = |r|$. For this monopole background, the charge distribution remains homogenous for any applied field.

We conclude that the coupling between itinerant electrons and local moment gives rise to an instability in the charge distribution in the presence of an applied electric field. The simplest steady state of this nonequilibrium problem has an inhomogenous charge density that, for a helical magnet, is a charge density wave of the same period as the helix. In general, magnetic configurations with constant magnitude but slowly varying direction give rise to charge accumulation only in an electric

field, while configurations with variable magnitude cause charge accumulation in equilibrium. However, there are

special configurations where charge homogeneity is protected for any an applied field.

-
- ¹ Q.Niu and T.Jungwirth, Phys. Rev. Lett. **88**, 207208 (2002).
 - ² F. D. M. Haldane, cond-mat/0408417 (2004).
 - ³ S. Murakami, N. Nagaosa, and S.-C. Zhang, Science **301**, 1348 (2003).
 - ⁴ Q.Niu and J.Sinova, Phys. Rev. Lett. **92**, 126603 (2004).
 - ⁵ S. Murakami, N. Nagosa, and S.-C. Zhang, Phys. Rev. B **69**, 214512 (2004).
 - ⁶ C. Pfleiderer, G. J. McMullan, S. R. Julian, and G. G. Lonzarich, Phys. Rev. B **55**, 8330 (1997).
 - ⁷ C. Pfleiderer, D. Reznik, L. Pintschovius, M. Garst, and A. Rosch, Nature **427**, 227 (2004).
 - ⁸ P. M. Woodward, D. E. Cox, E. Moshopoulou, A. W. Sleight, , and S. Morimoto, Phys. Rev. B **62**, 844 (2000).
 - ⁹ H. Ohno, J. Magn. Magn. Mater **200**, 110 (1999).
 - ¹⁰ H. J. Williams, J. H. Wernick, R. C. Sherwood, et al., J. Appl. Phys **37**, 1256 (1966).